

Divergence-free WKB method

Tadanori Hyouguchi¹, Satoshi Adachi¹, and Masahito Ueda^{1,2}

¹*Department of Physics, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8551, Japan*

²*CREST, Japan Science and Technology Corporation (JST), Saitama 332-0012, Japan*
(February 1, 2008)

A new semiclassical approach to linear (L) and nonlinear (NL) one-dimensional Schrödinger equation (SE) is presented. Unlike the usual WKB solution, our solution does not diverge at the classical turning point. For LSE, our zeroth-order solution, when expanded in powers of $\hbar$, agrees with the usual WKB solution $e^{iS/\hbar}/\sqrt{v}$ up to $\mathcal{O}(\hbar^0)$, where S and v are the classical action and the velocity, respectively. For NLSE, our zeroth-order solution includes quantum corrections to the Thomas-Fermi solution, thereby giving a smoothly decaying wave function into the forbidden region.

03.65.Ge, 03.65.Sq

The WKB method allows us to derive expressions for various quantum-mechanical quantities when the action is much larger than $\hbar$, and has been widely used in many subfields of physics and chemistry [1–4]. Nevertheless, the WKB method has a serious flaw of divergence at the classical turning point, which originates from the fact that the classical trajectory is taken as the zeroth-order solution. Considerable efforts have been devoted to overcome this problem, among other things, the complex method [1,2] and the uniform approximation [5,6]. However, the former method cannot describe the wave function at the turning point, while the latter one requires us to resort to a comparison function [7] that mimics the original potential and to which an analytic solution is available. This Letter presents a new semiclassical method whose zeroth-order solution is constructed upon a trajectory with quantum corrections such that the aforementioned divergence is absent. Moreover, our method gives a solution to the nonlinear Schrödinger equation (NLSE) on an equal footing.

We begin by reviewing the WKB method for the one-dimensional linear Schrödinger equation (LSE) $-\frac{\hbar^2}{2m}\frac{d^2\psi}{dx^2} + V\psi = E\psi$. Rescaling the length and the energy in units of l and $\hbar^2/2ml^2$, respectively, where l is a characteristic length scale of the potential $V(x)$, LSE takes the form

$$-\psi'' + V\psi = E\psi, \quad (1)$$

where the prime denotes the differentiation with respect to x . In Eq. (1), the length x and the energies E, V are proportional to $\hbar^0$ and $\hbar^{-2}$, respectively. We introduce the action function $\varphi(x)$ through $\psi(x) = e^{\varphi(x)}$, where the action is measured in units of $\hbar$. In terms of $\varphi(x)$, Eq. (1) takes the form

$$(\varphi')^2 + (E - V) = -\varphi'' . \quad (2)$$

The zeroth-order WKB solution $\varphi'_{\text{WKB},0}$, which is obtained by neglecting φ'' in Eq. (2), satisfies the following equation:

$$(\varphi'_{\text{WKB},0})^2 + (E - V) = 0 . \quad (3)$$

The classical turning point $x^{(c)}$ is determined by $V(x^{(c)}) = E$. Incorporating the effect of φ'' perturbatively, the WKB solution φ'_{WKB} takes familiar forms as

$$\varphi'_{\text{WKB}} = \begin{cases} \pm i \sqrt{E - V} - \frac{1}{4} \frac{V'}{V - E} + \cdots & (V < E), \\ \pm \sqrt{V - E} - \frac{1}{4} \frac{V'}{V - E} + \cdots & (V > E). \end{cases} \quad (4)$$

We note that the WKB solution diverges at $x^{(c)}$. This divergence originates from the fact that the classical trajectory is taken as the zeroth-order solution. Our strategy is to incorporate quantum corrections in the zeroth-order solution so as to remove the divergence.

To carry out this program, we differentiate both sides of Eq. (2) with respect to x , having

$$2\varphi'\varphi'' - V' = -\varphi''' , \quad (5)$$

and substitute φ'' in Eq. (2) into Eq. (5), obtaining

$$(\varphi')^3 + (E - V)\varphi' + \frac{1}{2}V' = \frac{1}{2}\varphi''' . \quad (6)$$

The Schrödinger equation (1) is sufficient for Eq. (6) to hold, but it is not necessary. To show this, we add $\varphi'\varphi''$ to both sides of Eq. (6) and integrate the resulting equation with respect to x . We then get

$$ge^{2\varphi} = (\varphi')^2 + (E - V) + \varphi'' , \quad (7)$$

where g is a constant of integration. Equation (7) can be rewritten as

$$-\psi'' + V\psi + g\psi^3 = E\psi . \quad (8)$$

This is nothing but NLSE which includes LSE as a particular case of $g = 0$. Throughout this Letter, we assume that for NLSE the wave function is normalized to unity $\int_{-\infty}^{\infty} dx |\psi(x)|^2 = 1$, while for LSE the normalization constant is left arbitrary.

Our zeroth-order solution φ'_0 , which is obtained by neglecting φ''' in Eq. (6), satisfies the following cubic equation

$$(\varphi'_0)^3 + 3p\varphi'_0 + 2q = 0 , \quad (9)$$

where $p \equiv (E - V)/3$ and $q \equiv V'/4$. Comparing Eqs. (6) and (9) with Eqs. (2) and (3), respectively, we see our method is a natural extension of the WKB method. Equation (9) is the fundamental equation of our new semiclassical method. As we shall see later, the solutions of Eq. (9) lead to divergence-free wave functions.

A root of the determinant of Eq. (9), $D \equiv p^3 + q^2 = 0$, determines our turning point $x = x^{(q)}$, which includes quantum corrections. In fact, the difference between $x^{(q)}$ and $x^{(c)}$ may be expanded as

$$x^{(q)} - x^{(c)} = \frac{3}{2} \frac{1}{[2V'(x^{(c)})]^{1/3}} \left(1 + \frac{\epsilon}{2} + \mathcal{O}(\epsilon^2, \delta) \right) , \quad (10)$$

where

$$\epsilon \equiv \frac{V''(x^{(c)})}{[2V'(x^{(c)})]^{4/3}} \quad \text{and} \quad \delta \equiv \frac{V'''(x^{(c)})}{[2V'(x^{(c)})]^{5/3}} .$$

We first discuss the zeroth-order solution. In the allowed region, the cubic equation (9) has the following three solutions:

$$\varphi'_0 = \begin{cases} -\kappa, \\ \frac{1}{2}\kappa + ik \equiv \varphi'_+ , \\ \frac{1}{2}\kappa - ik \equiv \varphi'_- , \end{cases} \quad (11)$$

where

$$\begin{cases} \kappa(x) \equiv & (q + \sqrt{D})^{1/3} + (q - \sqrt{D})^{1/3} \\ k(x) \equiv & \frac{\sqrt{3}}{2} \left[(q + \sqrt{D})^{1/3} - (q - \sqrt{D})^{1/3} \right] \end{cases} . \quad (12)$$

When expanded in powers of $\hbar$, Eq. (11) reduces to

$$\varphi'_0 = \begin{cases} -\kappa = & -\frac{2q}{3p} + \mathcal{O}(\hbar^1) = & \frac{1}{2} \frac{V'}{V-E} + \mathcal{O}(\hbar^1) , \\ \varphi'_+ = & i\sqrt{3p} + \frac{q}{3p} + \mathcal{O}(\hbar^1) = & i\sqrt{E-V} - \frac{1}{4} \frac{V'}{V-E} + \mathcal{O}(\hbar^1) , \\ \varphi'_- = & -i\sqrt{3p} + \frac{q}{3p} + \mathcal{O}(\hbar^1) = & -i\sqrt{E-V} - \frac{1}{4} \frac{V'}{V-E} + \mathcal{O}(\hbar^1) . \end{cases} \quad (13)$$

where $-\kappa$ and $\varphi'_\pm$, respectively, correspond to the Thomas-Fermi solution [8,9] and the WKB solutions (4) up to $\mathcal{O}(\hbar^0)$. Substituting Eq. (13) into Eq. (7), we obtain

$$g\psi^2 = \begin{cases} E - V + \mathcal{O}(\hbar^0) & \text{for } \varphi'_0 = -\kappa , \\ \mathcal{O}(\hbar^0) & \text{for } \varphi'_0 = \varphi'_+ , \\ \mathcal{O}(\hbar^0) & \text{for } \varphi'_0 = \varphi'_- . \end{cases} \quad (14)$$

From Eq. (14), it is clear that $-\kappa$ and $\varphi'_\pm$ give a solution to NLSE and those to LSE, respectively.

In the forbidden region, the cubic equation (9) has the following three real solutions :

$$\varphi'_0 = \begin{cases} -\kappa = \mp 2\sqrt{-p} \cos\left(\frac{1}{3}\text{Arctan}\frac{\sqrt{-D}}{|q|}\right), \\ \chi'_+ = \mp 2\sqrt{-p} \cos\left(\frac{1}{3}\text{Arctan}\frac{\sqrt{-D}}{|q|} + \frac{2\pi}{3}\right), \\ \chi'_- = \mp 2\sqrt{-p} \cos\left(\frac{1}{3}\text{Arctan}\frac{\sqrt{-D}}{|q|} - \frac{2\pi}{3}\right), \end{cases} \quad (15)$$

where the $-$ and $+$ signs correspond to $V' > 0$ and $V' < 0$, respectively. When expanded in powers of $\hbar$, Eq. (15) reduces to

$$\varphi'_0 = \begin{cases} -\kappa = \mp\sqrt{-3p} + \frac{q}{3p} + \mathcal{O}(\hbar^1) = \mp\sqrt{V-E} - \frac{1}{4}\frac{V'}{V-E} + \mathcal{O}(\hbar^1), \\ \chi'_+ = \pm\sqrt{-3p} + \frac{q}{3p} + \mathcal{O}(\hbar^1) = \pm\sqrt{V-E} - \frac{1}{4}\frac{V'}{V-E} + \mathcal{O}(\hbar^1), \\ \chi'_- = -\frac{2q}{3p} + \mathcal{O}(\hbar^1) = \frac{1}{2}\frac{V'}{V-E} + \mathcal{O}(\hbar^1), \end{cases} \quad (16)$$

where $-\kappa$ and χ'_+ correspond to the WKB solutions (4) up to $\mathcal{O}(\hbar^0)$. Substituting Eq. (16) into Eq. (7), we obtain

$$g\psi^2 = \begin{cases} \mathcal{O}(\hbar^0) & \text{for } \varphi'_0 = -\kappa, \\ \mathcal{O}(\hbar^0) & \text{for } \varphi'_0 = \chi'_+, \\ E - V + \mathcal{O}(\hbar^0) & \text{for } \varphi'_0 = \chi'_-. \end{cases} \quad (17)$$

To proceed further with our analysis, we shall assume that $V'(x) \geq 0$ ($\forall x$). Then the wave function must decay to zero as $x \rightarrow \infty$, *i.e.*, $\lim_{x \rightarrow \infty} \psi(x) = 0$. Consequently, in the forbidden region, $-\kappa$ must be chosen as the zeroth-order solution φ'_0 for both cases of LSE and NLSE. Note that, for NLSE, $g\psi^2 = \mathcal{O}(\hbar^0)$ in Eq. (17) does not mean that $g = 0$ but that the wave function is well damped in the forbidden region.

We are now in a position to construct our zeroth-order wave functions to LSE and NLSE. For NLSE, the zeroth-order wave function is

$$\psi_0(x) = N \exp\left(-\int_{x^{(q)}}^x dx' \kappa(x')\right), \quad (18)$$

where N is a normalization constant. Note that this single solution covers both allowed and forbidden regions.

For LSE, the zeroth-order wave function can be described, in general, as follows

$$\psi_0(x) = \begin{cases} A_+ \exp\left(\int_{x^{(q)}}^x dx' \varphi'_+(x')\right) + A_- \exp\left(\int_{x^{(q)}}^x dx' \varphi'_-(x')\right) \equiv \psi_{\text{I}}(x) & (x < x^{(q)}), \\ \exp\left(-\int_{x^{(q)}}^x dx' \kappa(x')\right) \equiv \psi_{\text{II}}(x) & (x > x^{(q)}). \end{cases} \quad (19)$$

To determine the relation between constants A_+ and A_- , we note that, near $x^{(c)}$, LSE (1) reduces to

$$-\frac{d^2\psi}{d\xi^2} + \xi\psi = 0, \quad (20)$$

where $\xi \equiv [V'(x^{(c)})]^{1/3}(x - x^{(c)})$. The exact solution of Eq. (20) that satisfies the boundary condition $\lim_{\xi \rightarrow \infty} \psi(\xi) = 0$ is the Airy function $\text{Ai}(\xi)$ [10]. In the region $|\xi| \gg 1$, the asymptotic forms of $\text{Ai}(\xi)$ are

$$\text{Ai}(\xi) \sim \begin{cases} \frac{1}{\sqrt{\pi}}(-\xi)^{-1/4} \sin\left(\frac{2}{3}(-\xi)^{3/2} + \frac{\pi}{4}\right) & (\xi < 0), \\ \frac{1}{2\sqrt{\pi}}\xi^{-1/4} \exp\left(-\frac{2}{3}\xi^{3/2}\right) & (\xi > 0). \end{cases} \quad (21)$$

On the other hand, for the linear potential, where $p = -[V'(x^{(c)})]^{2/3}\xi/3$ and $q = V'(x^{(c)})/4$, one can integrate $\varphi'_\pm$ and κ in Eq. (19); then the asymptotic behavior of Eq. (19) for $|\xi| \gg 1$ becomes

$$\psi_0 = \begin{cases} \psi_{\text{I}} \sim (2e)^{-2/3} 2^{1/3} (-\xi)^{-1/4} [A_+ \exp(-i(\frac{2}{3}(-\xi)^{3/2} + \frac{\pi}{4})) + A_- \exp(i(\frac{2}{3}(-\xi)^{3/2} + \frac{\pi}{4}))] & (\xi < 0), \\ \psi_{\text{II}} \sim \frac{1}{2}(2e)^{5/6} 2^{1/3} \xi^{-1/4} \exp(-\frac{2}{3}\xi^{3/2}) & (\xi > 0). \end{cases} \quad (22)$$

For Eq. (22) to match Eq. (21), we must choose $A_+/A_- = -1$. The zeroth-order solution (19) can thus be described as

$$\psi_0(x) = \begin{cases} \psi_{\text{I}} = A \exp\left(\frac{1}{2} \int_{x^{(\text{q})}}^x dx' \kappa(x')\right) \sin\left(\int_x^{x^{(\text{q})}} dx' k(x')\right) & (x < x_0), \\ \psi_{\text{II}} = \exp\left(-\int_{x^{(\text{q})}}^x dx' \kappa(x')\right) & (x > x_0), \end{cases} \quad (23)$$

where $A \equiv \mp 2iA_{\pm}$. For ψ_{I} and ψ_{II} and their first derivatives to be continuous at a certain point $x_0 (< x^{(\text{q})})$, *i.e.*, $\psi_{\text{I}}(x_0) = \psi_{\text{II}}(x_0)$ and $\psi'_{\text{I}}(x_0) = \psi'_{\text{II}}(x_0)$, x_0 and A must satisfy $\tan\left(\int_{x_0}^{x^{(\text{q})}} dx k(x)\right) = \frac{2k(x_0)}{3\kappa(x_0)}$ and $A = \exp\left(\frac{3}{2} \int_{x_0}^{x^{(\text{q})}} dx \kappa(x)\right) \text{cosec}\left(\int_{x_0}^{x^{(\text{q})}} dx k(x)\right)$. These simultaneous equations can be solved approximately, giving $x_0 = x^{(\text{c})} + \mathcal{O}(\epsilon)$ and $A = (2e)^{3/2} + \mathcal{O}(\epsilon)$. For the linear potential ($\epsilon = 0$), the asymptotic behavior of ψ_{I} (ψ_{II}) in Eq. (22) for $\xi \rightarrow -\infty$ ($+\infty$) with $A_{\pm} = \mp(2e)^{3/2}/2i$ exactly agrees with that of $\text{Ai}(\xi)$ in Eq. (21) except for an overall factor.

Our zeroth-order wave function (23) vanishes at $x^{(\text{q})}$ for any potential, which gives us a new quantization condition (compared with the WKB quantization condition)

$$\begin{cases} \frac{1}{2\pi} \oint dx k(x) = \frac{1}{\pi} \int_{x_{\text{L}}^{(\text{q})}}^{x_{\text{R}}^{(\text{q})}} dx k(x) = n + 1 & \text{(Our method),} \\ \frac{1}{2\pi} \oint dx k_{\text{WKB}}(x) = \frac{1}{\pi} \int_{x_{\text{L}}^{(\text{c})}}^{x_{\text{R}}^{(\text{c})}} dx k_{\text{WKB}}(x) = n + \frac{1}{2} & \text{(the WKB method),} \end{cases} \quad (24)$$

where $k(x)$ is given in Eq. (12), $k_{\text{WKB}}(x) \equiv \sqrt{E - V(x)}$, $n = 0, 1, 2, \dots$ is the number of nodes of the wave function, and the suffixes L and R denote the left- and right-side turning points, respectively. We note that $k(x)$ and $x^{(\text{q})}$ include quantum corrections, which accounts for an extra $1/2$ in our quantization condition.

We have so far ignored the term $\varphi'''/2$ on the right-hand side of Eq. (6). The effect of this term may be evaluated perturbatively, giving [11]

$$\varphi'_1 = \frac{1}{27} \left[-\varphi'_0 (V')^2 \frac{\left(\varphi'_0 - \frac{V''}{2V'}\right)^2}{((\varphi'_0)^2 + p)^4} + (V')^2 \frac{\varphi'_0 - \frac{V''}{2V'}}{((\varphi'_0)^2 + p)^3} + \frac{3}{2} V'' \frac{\varphi'_0 - \frac{V'''}{2V''}}{((\varphi'_0)^2 + p)^2} \right]. \quad (25)$$

The first-order wave function $\psi_1(x)$ can be constructed by using $\varphi'_0 + \varphi'_1$ instead of φ'_0 .

We finally apply the zeroth-order solution $\psi_0(x)$ and the first-order solution $\psi_1(x)$ to a linear potential (Figs. 1 and 2), a parabolic potential $V = x^2$ with $E = 17$ (Figs. 3 and 4), and the Morse potential $V = 900[\exp(-2x) - 2\exp(-x)]$ with $E = -(39/2)^2$ (Fig. 5). We also compare them with the exact solution and the WKB solution or a solution in which the Thomas-Fermi and WKB ones are combined.

As shown in the figures, the zeroth-order wave function ψ_0 does not diverge at the turning point but the error is discernible around the classical turning point $x^{(\text{c})}$. This small discrepancy is drastically improved by using the first-order wave function ψ_1 . The error associated with the zeroth-order solution is estimated to be [11]

$$\max \left| \frac{\psi_0 - \psi_{\text{exact}}}{\psi_{\text{exact}}} \right| \simeq 0.028 - 0.36\epsilon + \mathcal{O}(\epsilon^2, \delta). \quad (26)$$

Equation (26) implies that our zeroth-order wave functions (18) and (23) are suitable for $\epsilon > 0$ rather than $\epsilon < 0$; when $\epsilon < 0$, the wave function is unstable against a small perturbation.

In conclusion, we proposed a divergence-free semiclassical method that enables us to solve LSE and NLSE on an equal footing. Our zeroth-order solution is constructed upon a trajectory that includes quantum corrections and therefore allows a rapidly converging perturbative expansion of the wave function. Our method may therefore be applied to drastically improving related semiclassical methods such as instantons [12] and periodic orbital theory [13]. The results of these subjects will be reported elsewhere.

This work was supported by a Grant-in-Aid for Scientific Research (Grant No. 11216204) by the Ministry of Education, Science, Sports, and Culture of Japan, and by the Toray Science Foundation.

-
- [1] M. V. Berry and K. E. Mount, Rep. Prog. Phys. **35**, 315 (1972).
 - [2] N. Fröman and P. O. Fröman, *JWKB Approximation* (North-Holland, Amsterdam, 1965).
 - [3] W. H. Miller, Advan. Chem. Phys. **30**, 77 (1975).
 - [4] A. Voros, Ann. Inst. H. Poincaré. **39**, 211 (1983).
 - [5] R. E. Langer, Bull. Am. Math. Soc. **40**, 545 (1934); Phys. Rev. **51**, 669 (1937).
 - [6] S. C. Miller and R. H. Good, Phys. rev. **91**, 174 (1953).
 - [7] R. B. Dingle, Appl. Sci. Res. B **5**, 345 (1956).
 - [8] V. V. Goldman, I. F. Silvera and A. J. Leggett, Phys. Rev. B **24**, 2870 (1981).
 - [9] G. Baym and C. J. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
 - [10] *Handbook of Mathematical Functions* edited by M. Abramowitz and I. A. Stegun (Dover, New York, 1969).
 - [11] Detailed derivations will be described elsewhere.
 - [12] R. Rajarman, *Solitons and instantons* (North-Holland, Amsterdam, 1982).
 - [13] M. C. Gutzwiller, J. Math. Phys. **8**, 1979 (1967); **10**, 1004 (1969); **11**, 1791 (1970); **12**, 343 (1971).




FIG. 1. Zeroth-order solution ψ_0 (dashed curve) and first-order solution ψ_1 (dot-dashed curve) to LSE for a linear potential, $-d^2\psi/d\xi^2 + \xi\psi = 0$. The exact solution (solid curve) and the usual WKB solution (dotted curve) are superimposed for comparison. The region around the turning point is enlarged in the inset.




FIG. 2. Zeroth-order solution ψ_0 (dashed curve) and first-order solution ψ_1 (dot-dashed curve) to NLSE for a linear potential, $-d^2\psi/d\xi^2 + \xi\psi + \psi^3 = 0$. The exact solution (solid curve) and a solution in which the Thomas-Fermi and WKB ones are combined (dotted curve) are superimposed for comparison. The region around the turning point is enlarged in the inset.




FIG. 3. Solutions to LSE for a parabolic potential, $V = x^2$ with $E = 17$ and $\epsilon = 0.0476$. The notations are the same as FIG. 1. The region around the turning point is enlarged in the inset.




FIG. 4. Solutions to NLSE for a parabolic potential, $V = x^2$ with $E = 17$ and $\epsilon = 0.0476$. The notations are the same as FIG. 2. The region around the turning point is enlarged in the inset.




FIG. 5. Solutions to LSE for the Morse potential, $V = 900[\exp(-2x) - 2\exp(-x)]$ with $E = -(39/2)^2$, $\epsilon_L = 0.0982$ and $\epsilon_R = -0.0394$. The notations are the same as FIG. 1. The regions around the two turning points are enlarged in the insets.